Epilepsy Research, 17 (1994) 273–274
0920-1211/93/$06.00 © 1993 Elsevier Science Publishers B.V. All rights reserved

Author Index to Volume 17 (1994)

Abe, S., see Nakamura, M., 249
Ackermann, R.F., see Rocha, L., 135
Akazawa, K., see Nakamura, M., 249
Anderson, W.R., Franck, J.E., Stahl, W.L. and Maki, A.A., Na,K-ATPase is decreased in hippocampus of kainate-lesioned rats, 221
Applegate, C.D., see Gelbard, H.A., 23
Avoli, M., see Liu, Z., 237

Banerjee, P.K. and Snead III, O.C., Thalamic mediodorsal and intralaminar nuclear lesions disrupt the generation of experimentally induced generalized absence-like seizures in rats, 193
Bianchetti, A., see Cocito, L., 49
Blisard, K.S., Fagin, K., Falivena, P., Privitera, M.D., Olejniczak, P., Harrington, D.A., Taylor, K.L. and Scremin, O.U., Experimental seizures in the frog (*Rana pipiens*), 13
Bossi, L., see Cocito, L., 49
Burger, R.L., see Doretto, M.C., 157

Chishaki, A., see Nakamura, M., 249
Chugani, H.T., see Rocha, L., 135
Cocito, L., Bianchetti, A., Bossi, L., Giberti, L. and Loeb, C., GABA and phosphatidylserine in human photosensitivity: a pilot study, 49
Coenen, A.M.L., see Romijn, H.J., 31
Craggs, M., see Dürmüller, N., 167

Dailey, J.W., see Doretto, M.C., 157
Daniels, A.H., see Fujikawa, D.G., 207
Delgado, M.R., see Liu, H., 257
Desjardins, G.C., see Liu, Z., 237
de Waal, L.P., see van Engelen, B.G.M., 43
Doretto, M.C., Burger, R.L., Mishra, P.K., Garcia-Cairasco, N., Dailey, J.W.. and Jobe, P.C., A microdialysis study of amino acid concentrations in the extracellular fluid of the substantia nigra of freely behaving GEPR-9s: relationship to seizure predisposition, 157
Dürmüller, N., Craggs, M. and Meldrum, B.S., The effect of the non-NMDA receptor antagonists GYKI 52466 and NBQX and the competitive NMDA receptor antagonist D-CPPene on the development of amygdala kindling and on amygdala-kindled seizures, 167

Eadie, M.J., Epileptic seizures in 1902 patients: a perspective from a consultant neurological practice (1961–1991), 55
Engel Jr., J., see Rocha, L., 135

Fagin, K., see Blisard, K.S., 13
Falivena, P., see Blisard, K.S., 13
Franck, J.E., see Anderson, W.R., 221
Fujikawa, D.G., Daniels, A.H. and Kim, J.S., The competitive NMDA receptor antagonist CGP 40116 protects against status epilepticus-induced neuronal damage, 207

Garcia-Cairasco, N., see Doretto, M.C., 157
Gelbard, H.A. and Applegate, C.D., Persistent increases in dopamine D_2 receptor mRNA expression in basal ganglia following kindling, 23
Giberti, L., see Cocito, L., 49
Gilbert, M.E., The NMDA antagonist MK-801 suppresses behavioral seizures, augments afterdischarges, but does not block development of perforant path kindling, 145
Gloor, P., see Liu, Z., 237
Goto, Y., see Nakamura, M., 249

Harrington, D.A., see Blisard, K.S., 13

Iasemidis, L.D., Olson, L.D., Savit, R.S. and Sackellares, J.C., Time dependencies in the occurrences of epileptic seizures, 81

Jobe, P.C., see Doretto, M.C., 157

Kato, M., see Nakamura, M., 249
Khan, N. and Wieser, H.G., Limbic encephalitis: a case report, 175
Kim, J.S., see Fujikawa, D.G., 207

Liu, H. and Delgado, M.R., The influence of polytherapy on the relationships between serum carbamazepine and its metabolites in epileptic children, 257
Liu, Z., Nagao, T., Desjardins, G.C., Gloor, P. and Avoli, M., Quantitative evaluation of neuronal loss in the dorsal hippocampus in rats with long-term pilocarpine seizures, 237
Loeb, C., see Cocito, L., 49
Löscher, W. and Schmidt, D., Strategies in antiepileptic drug development: is rational drug design superior to random screening and structural variation?, 95

Maki, A.A., see Anderson, W.R., 221
Matsunaga, Y., see Mimaki, T., 233
McLean, M.J., see Wamil, A.W., 1
Meldrum, B.S., see Dürmüller, N., 167
Mimaki, T., Tanoue, H., Matsunaga, Y., Miyazaki, H. and Mino, M., Regional distribution of ^{14}C-zonisamide in rat brain, 233
Mino, M., see Mimaki, T., 233
Mishra, P.K., see Doretto, M.C., 157
Miyazaki, H., see Mimaki, T., 233

Nagao, T., see Liu, Z., 237
Nakamura, M., Abe, S., Goto, Y., Chishaki, A., Akazawa, K. and Kato, M., In vivo assessment of prevention of white-noise-induced seizure in magnesium-deficient rats by *N*-methyl-D-aspartate receptor blockers, 249

Olejniczak, P., see Blisard, K.S., 13
Olson, L.D., see Iasemidis, L.D., 81

Privitera, M.D., see Blisard, K.S., 13
Renier, W.O., see van Engelen, B.G.M., 43
Rocha, L., Ackermann, R.F., Chugani, H.T. and Engel Jr., J., Chronic pretreatment with naloxone modifies benzodiazepine receptor binding in amygdaloid kindled rats, 135
Romijn, H.J., Voskuyl, R.A. and Coenen, A.M.L., Hypoxic-ischemic encephalopathy sustained in early postnatal life may result in permanent epileptic activity and an altered cortical convulsive threshold in rat, 31

Sackellares, J.C., see Iasemidis, L.D., 81
Savit, R.S., see Iasemidis, L.D., 81
Schmidt, D., see Löscher, W., 95
Schwartzkroin, P.A., Cellular electrophysiology of human epilepsy, 185
Scremin, O.U., see Blisard, K.S., 13
Snead III, O.C., see Banerjee, P.K., 193
Stahl, W.L., see Anderson, W.R., 221

Tanoue, H., see Mimaki, T., 233
Taylor, K.L., see Blisard, K.S., 13

van Engelen, B.G.M., de Waal, L.P., Weemaes, C.M.R. and Renier, W.O., Serologic HLA typing in cryptogenic Lennox-Gastaut syndrome, 43
Voskuyl, R.A., see Romijn, H.J., 31

Wamil, A.W. and McLean, M.J., Limitation by gabapentin of high frequency action potential firing by mouse central neurons in cell culture, 1
Weemaes, C.M.R., see van Engelen, B.G.M., 43
Wieser, H.G., see Khan, N., 175